

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-EP/99-02

January 4, 1999

Distinction of atmospheric $\nu_\mu - \nu_\tau$ and $\nu_\mu - \nu_s$ oscillations using short or intermediate baseline experiments

A. Geiser ^{1 2}

Abstract

The current case for atmospheric ν_μ oscillations into active or sterile neutrinos is reviewed. It is argued that neither the study of neutral current events at Super-Kamiokande, nor the information obtained from future long baseline experiments might be sufficient to unambiguously decide between these two scenarios. However, a combination of these results with the results from future short or intermediate baseline τ appearance experiments would clearly resolve most of the remaining ambiguities. This conclusion does not strongly depend on whether the results from LSND will be confirmed or not. In the case that LSND would be confirmed, a negative result in such a short or intermediate baseline experiment would also unambiguously exclude the interpretation of LSND as indirect $\nu_\mu - \nu_\tau - \nu_e$ oscillations.

to be published in Eur. Phys. J. C

PACS codes: 14.60.Pq, 14.60.St

keywords: atmospheric, neutrino, nutau, sterile, oscillations, short, intermediate, baseline

¹CERN, EP Division, CH-1211 Genève 23, Tel. +41-22-7678564, Fax +41-22-7679070,
e-mail Achim.Geiser@cern.ch

²now at Lehrst. f. Exp. Physik IV, Dortmund University, Dortmund, Germany

1 Introduction

In the wake of the recent evidence for atmospheric ν_μ oscillations by Super-Kamiokande [1], one of the crucial issues is to clarify whether the observed effect is due to $\nu_\mu - \nu_\tau$, $\nu_\mu - \nu_e$, or $\nu_\mu - \nu_s$ (sterile) oscillations. The pure $\nu_\mu - \nu_e$ case is strongly disfavoured [2] by CHOOZ [3] and Super-Kamiokande [1], while pure $\nu_\mu - \nu_\tau$ and $\nu_\mu - \nu_s$ oscillations are equally possible [4]. Within the restrictions imposed by CHOOZ, complicated mixtures of the three cases can also not be excluded experimentally [5][6]. In fact, most models favouring the sterile neutrino interpretation [7][8][9][10] do suggest some (small) contribution from standard flavour oscillations. Due to the fundamental implications of the existence of a sterile neutrino for new physics, we will assume that a separation power of 3 to 5 σ is desired to rule out the mainly sterile oscillation scenario, and that a significance of at least 5 σ is required to establish it.

The paper is organized as follows: first, we will review the current experimental status of the active versus sterile oscillation hypothesis. We will then give some arguments why it is likely that future improvements on these measurements by atmospheric and long baseline neutrino experiments will leave important loopholes in the confirmation of either of the two hypotheses, and how a modest admixture of $\nu_\mu - \nu_\tau$ to mainly $\nu_\mu - \nu_s$ oscillations can mimic the pure $\nu_\mu - \nu_\tau$ oscillation case. Finally, we will show how these loopholes can be closed by using the (positive or negative) results from Short or Intermediate Baseline τ Appearance (SIBTA) experiments [11][12][13].

Furthermore, we will argue that this conclusion does not depend strongly on whether the LSND observation of $\nu_\mu - \nu_e$ oscillations [14] will be confirmed or not. If LSND would be confirmed, a negative result in a SIBTA experiment would also unambiguously exclude the interpretation of LSND as indirect $\nu_\mu - \nu_\tau - \nu_e$ oscillations, therefore ruling out all oscillation scenarios invoking this option [15]. This includes essentially all models trying to reconcile LSND with the atmospheric and solar neutrino result in the standard 3-neutrino framework [16].

2 Discussion of Super-Kamiokande indications

Results of neutrino oscillation experiments are often expressed in terms of an effective two flavour oscillation scheme with a mixing angle $\sin^2 2\theta$ between the two flavours and a mass difference δm^2 between the two relevant mass eigenstates. The most popular interpretation of the Super-Kamiokande results is to invoke maximal or close to maximal $\nu_\mu - \nu_\tau$ oscillations with a δm^2 in the $10^{-2} - 10^{-3} \text{ eV}^2$ range. Clearly, if this interpretation is correct, the low δm^2 excludes any observation of a $\nu_\mu - \nu_\tau$ oscillation signal in current [17][18] and future [11] short baseline experiments. However, a significant $\nu_\mu - \nu_e$ contribution to the angular dependence of the atmospheric neutrino result is not completely excluded, and even suggested by some of the models [16] trying to reconcile the atmospheric neutrino anomaly with LSND. In this case the constraint on $\delta m_{\mu\tau}^2$ could be considerably diluted, and τ appearance signals in short or intermediate baseline experiments could be possible.

Alternatively, several theoretical models suggest the interpretation of the Super-Kamiokande results as $\nu_\mu - \nu_s$ oscillations. In this case, the sterile neutrino could either be the right-handed (sterile) partner of the left-handed (active) muon neutrino (or its antiparticle), leading to maximal $\nu - \bar{\nu}$ oscillations analogous to $K^0 - \bar{K}^0$ oscillations [7][8], or the light

remnant of very massive neutrinos in GUT extensions of the standard model usually involving extra neutrino multiplets [9][10]. In most of these models the observation of $\nu_\mu - \nu_s$ oscillations could be further complicated by non-negligible admixtures of $\nu_\mu - \nu_\tau$ or $\nu_\mu - \nu_e$ oscillations.

Since, due to the high τ mass threshold, neither ν_τ nor ν_s produce a visible charged current (CC) signal in Super-Kamiokande, the two cases are experimentally almost indistinguishable in the standard Super-Kamiokande analysis of e-like and μ -like events [1][4]. However, ν_τ 's do produce neutral current (NC) interactions while sterile neutrinos do not. This could lead to a visible distinction in two kinds of measurements: In the $\nu_\mu - \nu_s$ case the up/down asymmetry observed in the CC sample should also be present for NC events, while no NC up/down asymmetry should occur in the $\nu_\mu - \nu_\tau$ (or $\nu_\mu - \nu_e$) case [19]. This effect also yields differences for the up/down asymmetry in inclusive event samples [20]. Furthermore, the NC suppression in the $\nu_\mu - \nu_s$ case modifies the predicted NC/CC ratios [21].

Unfortunately a clean NC/CC separation is experimentally difficult, and the expected effects are diluted by the (supposedly) unaltered contribution from ν_e NC events. The cleanest way to identify NC events in Super-Kamiokande is to require a single π^0 from the process $\nu + N \rightarrow \nu + N + \pi^0$, with N being either a neutron or a proton below Čerenkov threshold. The π^0 is detected via its decay into two photons which convert and yield two electron-like (double) rings whose invariant mass is consistent with the π^0 mass. This procedure reduces statistics by so much that currently no significant measurement of the up/down asymmetry can be obtained [22]. The ratio of two ring (π^0 -like, NC) to single ring (e-like, CC) events compared to the prediction for the no oscillation case is measured to be [22]³

$$\frac{(\pi^0/e)_{data}}{(\pi^0/e)_{pred}} = 0.88 \pm 0.08_{stat} \pm 0.19_{sys} \quad (1)$$

where the systematic error is dominated by the poorly known single π^0 cross section, and the statistical error is based on 535 days (2 years) of running. Assuming an initial ν_μ/ν_e ratio of 2.0, using the measured average ν_μ/ν_e ratio suppression of 0.63 [1], and assuming no background, the ratio in eq. (1) is expected to be 1.00 for $\nu_\mu - \nu_\tau$ oscillations (neither of the two contributions is affected), 0.75 for $\nu_\mu - \nu_s$ oscillations (the π^0 contribution is suppressed), 0.75 for $\nu_\mu - \nu_e$ oscillations (the e contribution is enhanced), and 0.94 for mainly $\nu_\mu - \nu_\tau$ oscillations with a 10% $\nu_\mu - \nu_e$ contribution. If there is significant background contamination these differences will be further reduced.

In order to disentangle $\nu_\mu - \nu_\tau$ from $\nu_\mu - \nu_s$ at the 3 σ level, the combined statistical and systematic error must therefore be reduced to 8% or less. If a 10% $\nu_\mu - \nu_e$ contribution is allowed, the required maximal uncertainty is reduced to 6%. Arbitrarily assuming 2140 days (8 years) of running, the statistical error will be about 4%. In order not to exceed the 6% total error this implies a systematic uncertainty of the order of 5% or less. Even with the planned measurement of the π^0 production cross section in the near detector of the K2K experiment [24] this seems to be very hard to achieve. We therefore conclude that this measurement will probably yield a useful indication, but is unlikely to firmly establish one of the two options.

Similar arguments hold for the π^0 up/down asymmetry. From a simple extrapolation of existing data [22], it looks unlikely that this method will distinguish the two cases by more

³The value of 0.94 quoted e.g. in [23] does not yet include background subtraction.

than about 2 standard deviations.

The possibility to separate ν_τ and ν_s from the measurement of the inclusive up/down asymmetry of multiring events, which also depends on the suppression of the NC contribution, is discussed in ref. [20]. Here, the statistical and systematic errors are smaller, but the differences are also small, again yielding a potential effect of about 2σ . Furthermore, $\nu_\mu - \nu_s$ oscillations with a significant $\nu_\mu - \nu_e$ admixture could yield the same asymmetry as pure $\nu_\mu - \nu_\tau$ oscillations.

3 Other atmospheric neutrino experiments

None of the currently planned atmospheric neutrino experiments [25][26][27][28] has a τ detection efficiency which is sufficiently large to see a significant pure charged current oscillation signal. In addition to the small cross section (τ mass threshold), the unknown direction of the incoming neutrino makes a significant kinematical analysis “à la NOMAD” [17] impossible. Emulsion techniques “à la CHORUS” [18] can not be used due to the inherently small target mass.

As in Super-Kamiokande, any efforts to distinguish between ν_s and ν_τ must therefore focus on neutral current events or on inclusive event rates. In a detector like NICE [26] there is a small window at $\delta m^2 \sim \text{few } 10^{-4} \text{ eV}^2$ where the oscillation pattern could actually be resolved, opening the possibility of comparing the energy dependence of the CC and NC event rates. In addition, such a low δm^2 would in itself be an indication for oscillations into active neutrinos, since the sterile case is somewhat disfavoured for such low δm^2 values due to earth matter effects which start to play a role [4].

Ref. [27] outlines an atmospheric neutrino detector concept which would allow to measure ν_τ appearance via an enhancement of muon-less events from τ decays at the highest accessible neutrino energies, where τ production is least suppressed. The nice feature of this concept is that ν_e events are effectively filtered out, therefore removing most of the ν_e background and the $\nu_\mu - \nu_\tau/\nu_\mu - \nu_e$ ambiguity. Combined with the NC suppression for ν_s this yields a $\nu_\mu - \nu_\tau/\nu_\mu - \nu_s$ separation of several standard deviations for $\delta m^2 \sim 5 \times 10^{-3} \text{ eV}^2$ or larger [27]. For $\delta m^2 \sim 3 \times 10^{-3}$ the sensitivity is significantly reduced (oscillations of high energy ν_μ 's are suppressed), while for even lower δm^2 the difference becomes marginal.

For the Super-Kamiokande most favoured case of $\delta m^2 \sim 2 \times 10^{-3} \text{ eV}^2$ it is not clear at present whether the difficulties outlined above will allow to draw any firm conclusions concerning the distinction of active and sterile neutrino oscillations. Furthermore, no detector of the types discussed above has been endorsed or approved so far.

In principle $\nu_\mu - \nu_\tau$ and $\nu_\mu - \nu_s$ oscillations can also be distinguished through the distortion of the momentum spectra of upward going muons due to matter effects [29]. No conclusion has been reached so far from ongoing experiments [30] due to large systematic errors. It is not clear at present whether these measurements can be improved sufficiently well to eventually allow a clear distinction.

4 Accelerator neutrino experiments

The best way to establish the $\nu_\mu - \nu_\tau$ interpretation of the atmospheric neutrino result obviously seems to be the detection of ν_τ appearance in long baseline accelerator experiments [28][31][32][33][25]. Here, appearance could be established either directly through the observation of τ production, or indirectly via an enhancement of the NC/CC ratio, together with the non-observation of a large electron appearance effect. But, it turns out that a positive τ signal in e.g. ICARUS [28], OPERA [31], or MINOS [32] would *not* automatically prove the $\nu_\mu - \nu_\tau$ hypothesis for atmospheric neutrinos, unless the corresponding δm^2 can be measured directly from this signal. The basic argument is that, as illustrated in Fig. 1, even a small $\nu_\mu - \nu_\tau$ contribution ($\sin^2 2\theta \sim \text{few } 10^{-3}$ or larger) at large δm^2 (order eV^2) can yield a signal in long baseline experiments (for both appearance and disappearance) that is similar in size or even larger than the expected effect from maximal $\nu_\mu - \nu_s$ oscillations, and can therefore mimic maximal $\nu_\mu - \nu_\tau$ oscillations at small δm^2 . This argument is discussed in detail below.

4.1 Motivation for mixed $\nu_\mu - \nu_s$ and $\nu_\mu - \nu_\tau$ scenarios

One of the most general arguments brought forward in favour of $\nu_\mu - \nu_s$ oscillations in the context of some models [7][9] is the possibility to combine a neutrino mass hierarchy and mixing pattern which are similar to the one observed in the quark and charged lepton sector (i.e. third generation is heaviest, mixing angles are small) with maximal $\nu_\mu - \nu_s$ oscillations, while retaining a significant contribution to hot dark matter. In mixed dark matter models, this would suggest a ν_τ mass of order eV. Similarity with the CKM matrix ([38]) would suggest a $\sin^2 2\theta_{\mu\tau}$ of order 10^{-2} (e.g. test point 2 in Fig. 1). Once sterile neutrinos are considered at all this is in some sense a “natural” possibility which should not be dismissed a priori, although it is by no means a necessity. Without requiring mass hierarchy, similar arguments for a possible $\nu_\mu \rightarrow \nu_\tau$ admixture apply for the models of ref. [10].

If LSND [14] would be confirmed this argument would be strengthened further, since the observation of $\nu_\mu - \nu_e$ oscillations in the range $\delta m^2 \sim 0.2 - 2 \text{ eV}^2$, combined with the hierarchy assumption, would again suggest a ν_τ mass in the eV range. In addition, it would make the sterile scenario more attractive, since the LSND result is hard to combine with the evidence from atmospheric and solar neutrinos in a three neutrino scheme. However, the argument given in the previous paragraph does *not* vanish if LSND would be disproven. Instead, $\nu_\mu - \nu_e$ oscillations might play a significant role at lower δm^2 , either as an admixture to mainly $\nu_\mu - \nu_s$ atmospheric neutrino oscillations or as the solution of the solar neutrino problem [39].

Finally, let us digress and consider the case of indirect $\nu_\mu - \nu_\tau - \nu_e$ oscillations in LSND [15][40]. Assuming mass hierarchy and denoting the dominant mass components of ν_e , ν_μ and ν_τ by m_1 , m_2 , and m_3 yields the mass relation $\delta m_{12}^2 \ll \delta m_{23}^2 \sim \delta m_{13}^2 \sim m_3^2$ known as one mass scale dominance [41]. In order for indirect oscillations to be detectable, m_3^2 must be in the range relevant for LSND, i.e. of order 1 eV^2 , and the rate proportional to

$$\sin^2 2\theta_{\text{LSND}} = 4|U_{e3}U_{\mu 3}|^2 \quad (2)$$

must be sufficiently large. Here, U_{e3} and $U_{\mu 3}$ are the relevant matrix elements of the general 3 neutrino mixing matrix. Bounds on U_{e3} from Bugey [42] combined with the LSND




Figure 1: Parameter space for $\nu_\mu - \nu_\tau$ oscillations. Indicated are the current combined limit from NOMAD [17], CHORUS [18], E531 [34] and CDHS [35] (thick continuous line) as well as the potential future limits from CHORUS/NOMAD, TOSCA [11], and a generic intermediate baseline experiment in the Jura [13] (dashed lines). The allowed regions of the Kamiokande (K) [36] and Super-Kamiokande (SK) [1] experiments, if interpreted as $\nu_\mu - \nu_\tau$ oscillations, are also shown. The bands labeled (b), (c), and (d) correspond to allowed regions from different potential τ appearance results of a generic long baseline experiment as discussed in the text, while (a) stands for a null result. The shaded area indicates the region favoured by mixed dark matter scenarios [37] while the points indicate test points used for the discussion in the text. They correspond to a specific prediction of $\nu_\mu - \nu_\tau$ oscillations (in addition to maximal $\nu_\mu - \nu_s$ oscillations) of ref. [8] (1), a generic test point compatible with most of the models in refs. [7][9][10] (2), and two test points corresponding to indirect oscillation solutions for LSND [15] (3 and 4). In addition, point 3 can be considered a variant of point 2.

measurement of the rate in eq. (2) yield a prediction for $U_{\mu 3}$ as a function of m_3^2 . All possible solutions are close to the limit from CDHS [35]. Two (marginally) allowed solutions, translated into $\sin^2 2\theta_{\mu\tau}$, are shown as test points 3 and 4 in Fig. 1.

Test point 3 corresponds to a scenario very similar to that of test point 2, where $\nu_\mu - \nu_\tau$ oscillations happen in addition to the $\nu_\mu - \nu_s$ oscillations responsible for the atmospheric neutrino deficit. It is therefore also relevant outside of the LSND context.

Test point 4 represents the region of very large $\nu_\mu - \nu_\tau$ mixing used, among others, for the Cardall/Fuller, Acker/Pakvasa and Thun/McKee schemes [16] without the need for sterile neutrinos.

4.2 Interpretation of long baseline observations

For the purpose of this study we will consider close to maximal ν_μ disappearance of atmospheric neutrinos to be an established fact, and anticipate that this will be confirmed⁴ through a positive effect in the ratio of the ν_μ CC rate in near and far detectors of the long baseline programme (K2K [24], MINOS [32], NICE [26], ...). Whenever atmospheric $\nu_\mu - \nu_\tau$ or $\nu_\mu - \nu_s$ oscillations are mentioned in the next few paragraphs, it is understood that there could be a small (up to 10%) contribution from $\nu_\mu - \nu_e$. A potentially even larger ν_e contribution is assumed to be measured and corrected for. Finally, Fig. 1 implies that very similar conclusions can be obtained from short and intermediate baseline experiments. We will therefore often refer to a generic Short or Intermediate Baseline τ Appearance (SIBTA) experiment in the discussion.

case (a): Long baseline experiments do not observe ν_τ appearance.

A positive signal in a SIBTA experiment would then *establish* that $\nu_\mu - \nu_\tau$ oscillations are outside of the range accessible to long baseline experiments, but within the range relevant to mixed dark matter (e.g. test point 1 in Fig. 1 for the short baseline case), and *force* the $\nu_\mu - \nu_s$ interpretation of the atmospheric neutrino result. The measured long baseline disappearance rate would yield a measurement of the δm^2 for $\nu_\mu - \nu_s$ oscillations, to be compared to the Super-Kamiokande result.

A negative result in a SIBTA experiment would *exclude* any $\nu_\mu - \nu_\tau$ contribution to the long baseline signal from the cosmologically relevant range. It would therefore *strongly suggest* the $\nu_\mu - \nu_\tau$ interpretation of the atmospheric neutrino oscillations at the low end of the Super-Kamiokande allowed δm^2 range, provided the observed long baseline disappearance signal (from a low energy beam) is consistent with this low δm^2 hypothesis.

case (b): Long baseline experiments observe a small ν_τ appearance signal.

Since the appearance signal is small (e.g. a handful of events in the case of direct τ appearance), it would supposedly *not* be possible to reliably measure the δm^2 from the energy distribution of the appearance signal alone. Again, a positive signal in a SIBTA experiment would *establish* that $\nu_\mu - \nu_\tau$ oscillations occur in the cosmologically relevant range (e.g. test point 2 in Fig. 1), and *force* the $\nu_\mu - \nu_s$ interpretation of the atmospheric neutrino result. The combination of the SIBTA and long baseline results would unambiguously fix the $\nu_\mu - \nu_\tau$ oscillation parameters. If the $\nu_\mu - \nu_s$ oscillations would occur at the Super-Kamiokande best fit point of $\delta m^2 \sim 2 \times 10^{-3} \text{ eV}^2$ (SK), they would be partially masked by the $\nu_\mu - \nu_\tau$ signal in

⁴But note some caveats explained later in the text.

the long baseline disappearance search. On the other hand, if the disappearance rate would turn out to be significantly larger than the appearance rate ($\delta m^2 > 2 \times 10^{-3} \text{ eV}^2$), the former could be used to measure the δm^2 of the $\nu_\mu - \nu_s$ oscillations.

A negative SIBTA result would again *exclude* any $\nu_\mu - \nu_\tau$ contribution to the long baseline signal from the cosmologically relevant range, and therefore *strongly suggest* the $\nu_\mu - \nu_\tau$ interpretation of the atmospheric neutrino oscillations with parameters close to the Super-Kamiokande best fit point. This can be cross-checked by requiring the appearance and disappearance rates to agree. The combination of the long baseline with the atmospheric neutrino results then yields a precise measurement of the $\nu_\mu - \nu_\tau$ oscillation parameters.

case (c): Long baseline experiments observe a large ν_τ appearance signal.

Given a large $\nu_\mu - \nu_\tau$ appearance signal, it might be possible to extract δm^2 from the energy distribution of the appearance signal alone, or at least to exclude a significant fraction of the available parameter space. However the example of the LSND experiment [14] shows that this possibility can not be taken for granted: Despite a claimed excess of 50 events over a small background, LSND is not able to differentiate the low and high δm^2 cases for their $\nu_\mu - \nu_e$ oscillation signal. Definitely, a low/high δm^2 distinction at the 5σ level would not be obvious for the long baseline results.

The preference of the low δm^2 case could e.g. suggest the $\nu_\mu - \nu_\tau$ interpretation of the atmospheric neutrino deficit with a δm^2 in the Kamiokande/Super-Kamiokande overlap region ($\sim 6 \times 10^{-2} \text{ eV}^2$). The requirement of *not* observing a SIBTA signal would essentially *eliminate* the whole large δm^2 range, including the regions suggested by dark matter and/or LSND (test point 3), and therefore strongly enhance the confidence in the low δm^2 result. Compatibility of the observed long baseline appearance and disappearance rates, although required, would *not* yield any further separation power, since a large τ appearance signal from e.g. test point 3 in Fig. 1 would completely mask the atmospheric oscillation effect, and also yield a consistent appearance/disappearance rate.

The preference of the high δm^2 case would imply the interpretation of the atmospheric neutrino anomaly as mainly $\nu_\mu - \nu_s$ oscillations. If LSND would be confirmed, it could in addition imply the compatibility of the $\nu_\mu - \nu_\tau$ oscillation parameters with the indirect oscillation hypothesis for LSND (test point 3). Given the importance of such a result, the cross-check from the requirement of a positive SIBTA result would be absolutely *essential*.

case (d): Long baseline experiments observe a close to maximal ($> 30\%$) ν_τ appearance signal.

Such a signal would be really spectacular, and inconsistent with the analysis of the Super-Kamiokande data in terms of two flavour oscillations. Since it would point to a δm^2 larger than 10^{-2} eV^2 it would either indicate a serious flaw in the Super-Kamiokande analysis, or require a three (or more) flavour scheme with significant contributions from two different δm^2 , one high and one low.

The energy distribution of the observed τ spectrum could already give a serious indication of the relevant high δm^2 . The rate of a positive SIBTA signal would however unambiguously fix the δm^2 for $\nu_\mu - \nu_\tau$ oscillations, and could decide whether it corresponds to the Cardall/Fuller or Thun/McKee solutions (test point 4) or to some lower δm^2 value.

A negative SIBTA result would *constrain* the δm^2 to about $2 \times 10^{-2} \text{ eV}^2$.

In either case the up/down asymmetry pattern of the Super-Kamiokande data suggesting

a significant low δm^2 contribution would either have to be disproven, attributed to $\nu_\mu - \nu_e$ oscillations through a corresponding observation in long baseline experiments, or attributed to a complicated mixture of $\nu_\mu - \nu_\tau$, $\nu_\mu - \nu_e$, and $\nu_\mu - \nu_s$ oscillations.

5 Compatibility with LSND

As outlined in the cases discussed above, none of the scenarios discussed essentially requires the confirmation of LSND. On the other hand, a confirmation of LSND would significantly enhance the interest in the $\nu_\mu - \nu_\tau / \nu_\mu - \nu_s$ distinction, since the addition of one or more sterile neutrinos might then be the only solution to simultaneously describe all the data. Also, the question whether the LSND signal is caused by direct or indirect oscillations becomes very relevant. As can be seen from Fig. 1, a negative SIBTA signal would unambiguously exclude the indirect oscillation possibility (test points 3 and 4), while a positive signal might allow it. A positive $\nu_\mu - \nu_\tau$ signal of kind (c) or (d) in the long baseline experiments could confirm this scenario, while a signal of kind (b) or (a) (no signal) would again exclude it. Finally, reversing the argument, the observation or non-observation of a signal in a SIBTA experiment before the LSND case is settled could, depending on the context of the results of other experiments, indirectly contribute to the LSND verification.

6 Conclusion

It has been shown that, whatever the outcome of future atmospheric and long baseline neutrino experiments, the complementary information from a short or intermediate baseline τ appearance experiment could be crucial for the unambiguous interpretation of the long baseline results. Such an experiment is therefore needed to distinguish cleanly between the $\nu_\mu - \nu_\tau$ and $\nu_\mu - \nu_s$ interpretations of the atmospheric neutrino anomaly, and even a negative result is very relevant in this context. In the absence of this cross-check, a small $\nu_\mu - \nu_\tau$ contribution at high δm^2 could mask the long baseline $\nu_\mu - \nu_s$ signal, therefore yielding a false confirmation of the interpretation of the atmospheric neutrino anomaly in terms of $\nu_\mu - \nu_\tau$ oscillations. This conclusion does not depend on whether LSND is confirmed or not, although a confirmation of LSND would enhance the interest in resolving this ambiguity. If LSND would be confirmed, a negative signal in such a short or intermediate baseline experiment would definitively rule out the indirect oscillation interpretation of the LSND result.

7 Acknowledgements

Constructive discussions with E. do Couto e Silva and J.J. Gomez-Cadenas are gratefully acknowledged. Special thanks go to the organizers of the NOW'98 neutrino workshop in Amsterdam, 6-9 September 1998, for stimulating the present work.

References

- [1] Super-Kamiokande Collaboration, Y. Fukuda et al., Phys. Rev. Lett **81** (1998) 1562;
Super-Kamiokande Collaboration, Y. Fukuda et al., Phys. Lett. **B 436** (1998) 33;
Super-Kamiokande Collaboration, Y. Fukuda et al., Phys. Lett. **B 433** (1998) 9.
- [2] M.C. Gonzalez-Garcia et al., Phys. Rev. **D58** (1998) 033004.
- [3] CHOOZ Collaboration, M. Apollonio et al., Phys. Lett. **B 420** (1998) 397.
- [4] R. Foot, R.R. Volkas, and O. Yasuda, Phys. Rev. **D58** (1998) 013006;
M.C. Gonzalez-Garcia et al., FTUV-98-56 and hep-ph/9807305, July 1998.
- [5] G.L. Fogli et al., Phys. Lett. **B 425** (1998) 341;
G.L. Fogli et al., BARI-TH-309-98 and hep-ph/9808205, Aug. 1998;
S.M. Bilenky and C. Giunti, DFTT 05/98 and hep-ph/9802201, Feb. 1998;
R. Foot, R.R. Volkas, and O. Yasuda, Phys. Lett. **B 433** (1998) 82.
- [6] V. Barger et al., Phys. Rev. **D 58** (1998) 093016.
- [7] A. Geiser, CERN-EP-98-056, Apr. 1998, submitted to Phys. Lett. **B**;
M. Kobayashi, C.S. Lim and M.M. Nojiri, Phys. Rev. Lett. **67** (1991) 1685;
C. Giunti, C.W. Kim and U.W. Lee, Phys. Rev. **D 46** (1992) 3034.
- [8] Y. Koide and H. Fusaoka, US-98-06 and hep-ph/9806516, June 1998;
Y. Koide, Phys. Rev. **D 57** (1998) 5836.
- [9] J.P. Bowes and R.R. Volkas, J. Phys. **G 24** (1998) 1249;
R. Foot and R.R. Volkas, Phys. Rev. **D 52** (1995) 6595;
R. Foot, Mod. Phys. Lett. **A 9** (1994) 169.
- [10] Q.Y. Liu and A.Yu. Smirnov, Nucl. Phys. **B 524** (1998) 505;
Q.Y. Liu and A.Yu. Smirnov, IC-98-14, Febr. 1998;
E.J. Chun, C.W. Kim, and U.W. Lee, Phys. Rev. **D 58** (1998) 093003;
A.S. Joshipura and A.Yu. Smirnov, PRL-TH-98-005 and hep-ph/9806376, June 1998;
M. Bando and K. Yoshioka, KUNS-1516 and hep-ph/9806400, June 1998.
- [11] TOSCA Letter of Intent, CERN-SPSC/97-5, SPSC/I 213 (1997);
Addendum to Letter of Intent, CERN-SPSC/98-20, SPSC/I 213 add., May 1998.
- [12] R. A. Sidwell (COSMOS Collaboration) in Neutrino '96, Proceedings of the 17th International Conference on Neutrino Physics and Astrophysics, Helsinki, Finland, June 13, 1996, ed. by K. Enqvist, K. Huitu and J. Maalampi, World Scientific (1997), p. 152.
- [13] proposal for ICARUS in the Jura, CERN/SPSLC 96-58 (1996);
D. Autiero et al., Expression of Interest for Jura experiment, CERN/SPSC/97-23, SPSC/I 217, Oct. 1997.
- [14] LSND Collaboration, C. Athanassopoulos et al., Phys. Rev. Lett. **75** (1995) 2650;
J.E. Hill, Phys. Rev. Lett. **75** (1995) 2654;
LSND Collaboration, C. Athanassopoulos et al., Phys. Rev. **C 54** (1996) 2685;

- LSND Collaboration, C. Athanassopoulos et al., Phys. Rev. Lett. **77** (1996) 3082;
 LSND Collaboration, C. Athanassopoulos et al., LA-UR-97-1998, UCRHEP-E191, and
 nucl-ex/9706006, June 1997, submitted to Phys. Rev. **C**;
 LSND Collaboration, C. Athanassopoulos et al., Phys. Rev. Lett. **81** (1998) 1774.
- [15] K.S. Babu, J.C. Pati, and F. Wilczek, Phys. Lett. **B 359** (1995) 351.
- [16] C.Y. Cardall and G.M. Fuller, Phys. Rev. **D 53** (1996) 4421;
 G.L. Fogli et al., Phys. Rev. **D56** (1997) 4365;
 A. Acker and S. Pakvasa, Phys. Lett. **B 397** (1997) 209;
 T. Teshima, T. Sakai and O. Inagaki, CU-TP-98-01 and hep-ph/9801276, Jan. 1998;
 R.P. Thun and S. McKee, UM-HE-98-11 and hep-ph/9806534, June 1998.
- [17] NOMAD Collaboration, J. Altegoer et al., Phys. Lett. **B 431** (1998) 219.;
 D. Autiero (NOMAD Collaboration), talk at ICHEP98 Vancouver, Canada, July 23-29
 1998.
- [18] CHORUS Collaboration, E. Eskut et al., Phys. Lett. **B 424** (1998) 202;
 CHORUS Collaboration, E. Eskut et al., CERN-EP/98-73;
 P. Migliozi (CHORUS Collaboration), talk at ICHEP98 Vancouver, Canada, July 23-29
 1998.
- [19] J.W. Flanagan, J.G. Learned, and S. Pakvasa, Phys. Rev. **D 57** (1998) 2649;
 J.G. Learned, S. Pakvasa, and J.L. Stone, Phys. Lett. **B 435** (1998) 131.
- [20] L.J. Hall, H. Murayama, Phys. Lett. **B 436** (1998) 323.
- [21] F. Vissani and A. Yu. Smirnov, Phys. Lett. **B 432** (1998) 376.
- [22] W. Gajewski (Super-Kamiokande Collaboration), talk at Neutrino Oscillation Work-
 shop (NOW 98), Amsterdam, Netherlands, 7-9 September 1998.
- [23] T. Kajita (Super-Kamiokande Collaboration), talk at 18th International Conference on
 Neutrino Physics and Astrophysics, Takayama, Japan, June 1998.
- [24] Y. Suzuki (E362 (K2K) Collaboration), in Neutrino '96 [12], p. 237.
- [25] NOE Collaboration, M. Ambrosio et al., Nucl. Instr. Meth. **A 363** (1995) 604.
- [26] NICE proposal, G. Baldini et al., LNGS-LOI 98/13 (1998)
- [27] A. Curioni et al., ICGF-98-05 and hep-ph/9805249, May 1998.
- [28] ICARUS Collaboration, P. Benetti et al., Nucl. Instr. Methods **A 327** (1993) 327; **A**
332 (1993) 332; P. Cennini et al., Nucl. Instr. Methods **A 333** (1993) 567; **A 345**
 (1994) 230; **A 355** (1995) 355;
 ICARUS-CERN-Milano Collaboration, CERN/SPSLC 96-58, SPSLC/P 304, December
 1996.

- [29] P. Lipari and M. Lusignoli, Phys. Rev. **D 58** (1998) 073005;
Q.Y. Liu, S.P. Mikheyev, and A.Y. Smirnov, IC/98/30 and hep-ph/9803415, March 1998, submitted to Phys. Lett. **B**.
- [30] MACRO collaboration, M. Ambrosio et al., Phys. Lett. **B 434** (1998) 451;
MACRO Collaboration, S. Ahlen et al., Phys. Lett. **B 357** (1995) 481;
Kamiokande Collaboration, S. Hatakeyama et al., Phys. Rev. Lett. **81** (1998) 2016;
Baksan Collaboration, review by T. Gaisser, in Neutrino '96 [12], p. 211;
IMB Collaboration, R. Becker-Szendy et al., Phys. Rev. **D 46** (1992) 3720.
- [31] OPERA letter of intent, S. Shibuya et al., CERN-SPSC/97-24, SPSC/I 218, LNGS-LOI-8-97 (1997).
- [32] MINOS Collaboration, Fermilab proposal P875;
MINOS Collaboration, "Neutrino Oscillation Physics at Fermilab: The NuMI-MINOS Project", NuMI-L375, May 1998;
S.G. Wojcicki, in Neutrino '96 [12], p. 231.
- [33] AquaRICH Collaboration, T. Ypsilantis et al., Nucl. Instr. Meth. **A 371** (1996) 330.
- [34] E531 Collaboration, N. Ushida et al., Phys. Rev. Lett. **57** (1986) 2897.
- [35] CDHS Collaboration, F. Dydak et al., Phys. Lett. **B 134** (1984) 281.
- [36] Kamiokande Collaboration, Y. Fukuda et al., Phys. Lett. **B 335** (1994) 237.
- [37] see e.g. J. Primack and A. Klypin, Nucl. Phys. Proc. Suppl. **51 B** (1996) 30.
- [38] N. Cabibbo, Phys. Rev. Lett **10** (1963) 531;
M. Kobayashi and T. Maskawa, Progr. Th. Phys. **49** (1973) 652;
F.J. Gilman, K. Kleinknecht and B. Renk, in Review of Particle Physics, Phys. Rev. **D 54** (1996), p. 94.
- [39] N. Hata and P. Langacker, Phys. Rev. **D 56** (1997) 6107.
- [40] C.Y. Cardall, G.M. Fuller, and D.B. Cline, Phys. Lett. **B 413** (1997) 246.
- [41] G.L. Fogli, E. Lisi, and G. Sciosia, Phys. Rev. **D 52** (1995) 5334.
- [42] Bugey Collaboration, B. Achkar et al., Nucl. Phys. **B 434** (1995) 503.